

AD-A147 524

A PROPAGATOR EXPANSION METHOD FOR SOLVING LINEARIZED  
PLASMA KINETIC EQUATIONS WITH COLLISIONS(U) AIR FORCE  
GEOPHYSICS LAB HANSCOM AFB MA J R JASPERSE 25 JUN 84

1/1


UNCLASSIFIED

AFGL-TR-84-0170

F/G 20/9

NL





12

AD-A147 524

AFGL-TR-84-0170  
ENVIRONMENTAL RESEARCH PAPERS, NO. 884

# A Propagator Expansion Method for Solving Linearized Plasma Kinetic Equations With Collisions

JOHN R. JASPERSE



25 June 1984



Approved for public release; distribution unlimited.

DTIC FILE COPY



DTIC  
ELECTE  
NOV 14 1984  
S E D

IONOSPHERIC PHYSICS DIVISION

PROJECT 2310

**AIR FORCE GEOPHYSICS LABORATORY**


HANSCOM AFB, MA 01731

84 11 05 085

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Information Service (NTIS).

"This technical report has been reviewed and is approved for publication."

FOR THE COMMANDER

  
\_\_\_\_\_  
Herbert C. Carlson, Jr.  
Branch Chief

  
\_\_\_\_\_  
Robert A. Skrivanek, Director  
Division Director

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify AFGL/DAA, Hanscom AFB, MA 01731. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS None		
2a. SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE N/A				
4. PERFORMING ORGANIZATION REPORT NUMBER(S) AFGL-TR-84-0170 ERP, No. 884		5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Air Force Geophysics Laboratory		6b. OFFICE SYMBOL (If applicable) LIS		7a. NAME OF MONITORING ORGANIZATION
6c. ADDRESS (City, State and ZIP Code) Hanscom AFB Massachusetts 01731		7b. ADDRESS (City, State and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER
8c. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FUNDING NOS.		
		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
				WORK UNIT NO.
11. TITLE (Include Security Classification) A Propagator Expansion Method for (over)		61102F	2310	G6 02
12. PERSONAL AUTHOR(S) John R. Jasperse				
13a. TYPE OF REPORT Scientific	13b. TIME COVERED FROM June 83 TO June 84	14. DATE OF REPORT (Yr., Mo., Day) 1984 June 25		15. PAGE COUNT 15
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES				
FIELD	GROUP	SUB. GR.		
20	09.14			
18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Operator expansion method, plasma kinetic equations, collisional plasmas				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) → In this paper, we present a propagator expansion method for solving linearized plasma kinetic equations with collisions. The essence of the method is the representation and use of the collisional propagator for a given problem as an expansion in powers of the collision frequency. For the linearized Balescu-Lenard collision operator and for the zero-order distribution function Maxwellian, we obtain an exact expression for the collisional damping rate in the long wavelength approximation to the first order in the plasma parameter. K				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL John R. Jasperse		22b. TELEPHONE NUMBER (Include Area Code) (617)861-3137		22c. OFFICE SYMBOL LIS

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

11. Title (Contd)

Solving Linearized Plasma Kinetic Equations With Collisions

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

## Preface

I should like to acknowledge important discussions with B. Basu, T.S. Chang, and J. M. Retterer. This work was sponsored by the Air Force Office of Scientific Research.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## Contents

1. INTRODUCTION	7
2. KINETIC EQUATIONS	8
3. EQUATION FOR THE COLLISIONAL PROPAGATOR	10
4. ASYMPTOTIC EXPANSIONS AS $\nu \rightarrow 0$	11
5. UNIFORMLY CONVERGENT EXPANSIONS IN POWERS OF $\nu$	12
6. SYMMETRY PROPERTIES	13
7. SOME APPLICATIONS	13
8. DISCUSSION	14
REFERENCES	15



# A Propagator Expansion Method For Solving Linearized Plasma Kinetic Equations With Collisions

## 1. INTRODUCTION

In 1946, Landau<sup>1</sup> gave a method for solving linearized plasma kinetic problems where discrete particle interactions were neglected. Subsequent studies<sup>2-4</sup> of collisional plasmas have employed methods of solution tailored to a specific form for the collision operator. In this paper, we give a general expansion method for solving linearized plasma kinetic problems when collisions are included. The method can be applied to a wide class of collision operators and has, for example, produced closed-form results for the collisional dielectric function for the Balescu<sup>5</sup>-Lenard<sup>6</sup> collision operator.

The essence of the idea presented in this paper is the derivation of a collisional propagator in terms of the collisionless propagator for the Vlasov equation and the linearized collision operator, and the representation and use

(Received for publication 21 June 1984)

1. Landau, L.D. (1946) Zh. Eksp. Teor. Fiz. 16:574 and J. Phys. (USSR) 10:25.
2. Bhatnagar, P.L., Gross, E.P., and Krook, M. (1954) Phys. Rev. 94:511.
3. Lenard, A., and Bernstein, I.B. (1958) Phys. Rev. 112:1456.
4. Dougherty, J.P. (1964) Phys. Fluids 7:1788.
5. Balescu, R. (1960) Phys. Fluids 3:52.
6. Lenard, A. (1960) Ann. Phys. (N.Y.) 3:390.

of the collisional propagator either as a series solution in powers of the collision frequency  $\nu$  or as an asymptotic expansion as the collision frequency tends to zero.

## 2. KINETIC EQUATIONS

The kinetic equations in the electrostatic approximation for a collisional plasma are:

$$\left\{ \frac{\partial}{\partial t} + \vec{v} \cdot \frac{\partial}{\partial \vec{r}} + \left[ \frac{q_e}{m_e} \vec{E}(\vec{r}, t) + \vec{a}_0(\vec{v}) \right] \cdot \frac{\partial}{\partial \vec{v}} \right\} f(\vec{r}, \vec{v}, t) = \mathcal{L}(f) , \quad (1)$$

$$\frac{\partial}{\partial \vec{r}} \cdot \vec{E}(\vec{r}, t) = 4\pi \left[ n_i q_i + q_e \int d^3v f(\vec{r}, \vec{v}, t) \right] , \quad (2)$$

where  $\vec{E}$  is the self-consistent, electrostatic field;  $\vec{a}_0$  is the force-per-unit mass due to spatially uniform, stationary fields; and  $\mathcal{L}(f)$  is the nonlinear collision operator that acts on the velocity-space coordinates of the electron distribution function  $f$ . The ions are assumed to provide a fixed neutralizing background with charge density  $n_i q_i$  while the other quantities have their usual meanings. We seek a perturbation solution of the form

$$f(\vec{r}, \vec{v}, t) = f_0(\vec{v}) + f_1(\vec{r}, \vec{v}, t) , \quad (3)$$

$$\vec{E}(\vec{r}, t) = \vec{E}_1(\vec{r}, t) , \quad (4)$$

where  $f_0$  is the zero-order solution and  $f_1$  and  $\vec{E}_1$  are perturbed quantities. The first-order equations using abbreviated notation are:

$$\left( \frac{\partial}{\partial t} + \vec{v} \cdot \frac{\partial}{\partial \vec{r}} + \vec{a}_0 \cdot \frac{\partial}{\partial \vec{v}} - L^\nu \right) f^\nu(t) = S^\nu(t) , \quad (5)$$

$$\frac{\partial}{\partial \vec{r}} \cdot \frac{\partial}{\partial \vec{r}} \phi^\nu(t) = -4\pi q_e \int d^3v f^\nu(t) , \quad (6)$$

$$S^\nu(t) = \frac{q_e}{m_e} \frac{\partial}{\partial \vec{r}} \cdot \phi^\nu(t) \cdot \frac{\partial}{\partial \vec{v}} f_0 , \quad (7)$$

where the subscript 1 on the perturbed quantities has been deleted and the superscript  $\nu$  is used to denote collisional dependence,  $\phi^\nu(t)$  is the electrostatic potential, and  $L^\nu$  denotes the linearized collision operator. The operator equation for the collisional propagator operator  $U^\nu(t-t')$ , is

$$\left( \frac{\partial}{\partial t} + \vec{v} \cdot \frac{\partial}{\partial \vec{r}} + \vec{a}_0 \cdot \frac{\partial}{\partial \vec{v}} - L^\nu \right) U^\nu(t-t') = 0 , \quad (8)$$

where  $U^\nu(t-t')$  is given by

$$U^\nu(t-t') = \int d^3r' \int d^3v' G^\nu(\vec{r}-\vec{r}', \vec{v}, \vec{v}', t-t') , \quad (9)$$

and where  $G^\nu(\vec{r}-\vec{r}', \vec{v}, \vec{v}', t-t')$ , the Green's function for the problem, is translationally invariant in space and time. Here,  $U^\nu(\tau)$  is subject to the causality condition that  $U^\nu(\tau)=0$  for  $\tau < 0$  and the initial condition that  $U^\nu(0)=1$ . In terms of  $U^\nu(t-t')$  the solution for  $f^\nu(t)$  is

$$f^\nu(t) = U^\nu(t) f^\nu(0) + \int_0^t dt' U^\nu(t-t') S^\nu(t') , \quad (10)$$

where  $f^\nu(0)$  denotes the initial value of  $f^\nu(t)$ . It is convenient to transform Eqs. (6) through (10) to  $\vec{k}-\vec{v}-\omega$  space using

$$f^\nu(t) = T_{\vec{k}, \omega}^{-1} \tilde{f}^\nu(\vec{k}, \vec{v}, \omega) = T_{\vec{k}, \omega}^{-1} \tilde{f}^\nu(\omega) , \quad (11)$$

$$\tilde{f}^\nu(\omega) = T_{\vec{r}, t}^{-1} f^\nu(\vec{r}, \vec{v}, t) = T_{\vec{r}, t}^{-1} f^\nu(t) , \quad (12)$$

$$T_{\vec{k}, \omega}^{-1} = \int \frac{d^3k}{(2\pi)^3} \int_C \frac{d\omega}{2\pi} \exp(i\vec{k} \cdot \vec{r} - i\omega t) , \quad (13)$$

$$T_{\vec{r}, t}^{-1} = \int d^3r \int_0^\infty dt \exp(-i\vec{k} \cdot \vec{r} + i\omega t) , \quad (14)$$

where  $\vec{k}$  is real,  $\text{Im } \omega \geq b > 0$ , and the contour  $C$  runs from  $-\infty + ib$  to  $+\infty + ib$ . Here  $b$  is chosen sufficiently large enough to ensure that the  $t$ -integration converges.<sup>7</sup> The equations in  $\vec{k}-\vec{v}-\omega$  space are:

$$\tilde{f}^\nu(\omega) = \tilde{U}^\nu(\omega) \left[ f_{\vec{k}}^\nu(0) + i \frac{q_e}{m_e} \cdot \tilde{\phi}^\nu(\omega) \vec{k} \cdot \frac{\partial}{\partial \vec{v}} f_0 \right] , \quad (15)$$

7. Titchmarsh, E. C. (1948) Introduction to the Theory of Fourier Integrals, Oxford University Press, London, Chapter 1.

$$\tilde{\phi}^{\nu}(\omega) = \frac{4\pi q_e}{k^2} \left[ \frac{1}{\epsilon^{\nu}(\omega)} \right] \int d^3v \tilde{U}^{\nu}(\omega) f^{\nu}_{\vec{k}}(0) , \quad (16)$$

$$\epsilon^{\nu}(\omega) = 1 - i \left( \frac{4\pi q_e^2}{m_e k^2} \right) \int d^3v \tilde{U}^{\nu}(\omega) \vec{k} \cdot \frac{\partial}{\partial \vec{v}} f_0 , \quad (17)$$

where  $f^{\nu}_{\vec{k}}(0)$  is the spatial transform of  $f^{\nu}(0)$ . The operator  $\tilde{U}^{\nu}(\omega)$  is

$$\tilde{U}^{\nu}(\omega) = \int d^3v' \tilde{G}^{\nu}(\vec{k}, \vec{v}, \vec{v}', \omega) , \quad (18)$$

and the Green's function in  $\vec{k}-\vec{v}-\omega$  space is

$$\tilde{G}^{\nu}(\vec{k}, \vec{v}, \vec{v}', \omega) = T_{\vec{R}, \tau} G^{\nu}(\vec{R}, \vec{v}, \vec{v}', \tau) . \quad (19)$$

### 3. EQUATION FOR THE COLLISIONAL PROPAGATOR

In order to derive an equation for  $\tilde{U}^{\nu}(\omega)$ , we consider a collisionless propagator operator,  $U(t-t')$ , which obeys the operator equation

$$\left( \frac{\partial}{\partial t} + \vec{v} \cdot \frac{\partial}{\partial \vec{r}} + \vec{a}_0 \cdot \frac{\partial}{\partial \vec{v}} \right) U(t-t') = 0 \quad (20)$$

and is subject to the same causality and initial conditions as those imposed on  $U^{\nu}(\tau)$ . We return to Eq. (8) and write an operator equation for  $U^{\nu}(t-t')$  in terms of  $U(t-t')$  and  $L^{\nu}$ :

$$U^{\nu}(t-t') = U(t-t') + \int_{t'}^t dt'' U(t-t'') L^{\nu} U^{\nu}(t''-t') , \quad (21)$$

where we have used causality and the initial conditions. Transforming Eq. (21) to  $\vec{k}-\vec{v}-\omega$  space, we obtain

$$\tilde{U}^{\nu}(\omega) = \tilde{U}(\omega) + \tilde{U}(\omega) L^{\nu} \tilde{U}^{\nu}(\omega) . \quad (22)$$

Solving Eq. (22) formally, we obtain

$$\tilde{U}^{\nu}(\omega) = [1 - \tilde{U}(\omega) L^{\nu}]^{-1} \tilde{U}(\omega) . \quad (23)$$

Equations (15) through (17) and Eq. (23) give a formal solution to the collisional, initial-value problem.

#### 4. ASYMPTOTIC EXPANSIONS AS $\nu \rightarrow 0$

In order to pursue a solution by the expansion method, we introduce a collision frequency  $\nu$  and write  $L^\nu = \nu L$ . In most cases,  $\nu$  will be small compared to other frequencies of interest. For a wide class of collision operators including integral, differential, and integral-differential, and for some continuity and integrability conditions on the functions that appear in the problem, it can be shown that the iteration of Eq. (22) yields an asymptotic expansion for  $\tilde{U}^\nu(\omega)$  as  $\nu \rightarrow 0$ :

$$\begin{aligned} \tilde{U}^\nu(\omega) \sim & \tilde{U}(\omega) + \nu \tilde{U}(\omega) L \tilde{U}(\omega) + \dots \\ & + \nu^n [\tilde{U}(\omega) L]^n \tilde{U}(\omega) + \dots \end{aligned} \quad (24)$$

By this we mean that if  $\tilde{g}(\omega) = \tilde{g}(\vec{k}, \vec{v}, \omega)$  and  $\tilde{f}^\nu(\omega) = \tilde{U}^\nu(\omega) \tilde{g}(\omega)$ , then the action of the iterated  $\tilde{U}^\nu(\omega)$ , given by Eq. (24), on  $\tilde{g}(\omega)$  generates an asymptotic expansion for  $\tilde{f}^\nu(\omega)$  as  $\nu \rightarrow 0$ . Similar results can be obtained for the iterated  $\epsilon^\nu(\omega)$ . If the iterated  $\tilde{f}^\nu(\omega)$  is a divergent asymptotic expansion, then Eq. (24) may not be truncated and used to approximate the initial-value problem as  $t \rightarrow \infty$  because time secularities will appear in the solution when transformed to  $\vec{r}-\vec{v}-t$  space. However, even if the iterated  $\epsilon^\nu(\omega)$  is also divergent but  $\nu/\omega_p \ll 1$ , where  $\omega_p$  is the plasma frequency, then  $\epsilon^\nu(\omega)$  may be truncated at first order in  $\nu$  and set equal to zero to obtain an approximate solution for the roots of the dispersion relation. This provides a solution for the normal mode and stability problems accurate to order  $\nu/\omega_p$ . The expression for the collisional dielectric function truncated to the first order in  $\nu$ , denoted by  $\epsilon^{(1)}(\vec{k}, \omega)$ , is

$$\epsilon^{(1)}(\vec{k}, \omega) = 1 - i \frac{\omega_p^2}{k^2} \{ T_0(\vec{k}, \omega) + \nu T_1(\vec{k}, \omega) \} , \quad (25)$$

$$T_n(\vec{k}, \omega) = \int d^3v [\tilde{U}(\omega) L]^n \tilde{U}(\omega) \vec{k} \cdot \frac{\partial}{\partial \vec{v}} f_0(\vec{v}) , \quad (26)$$

where  $f_0$  has been normalized to one.

## 5. UNIFORMLY CONVERGENT EXPANSIONS IN POWERS OF $\nu$

For the special case when  $\hat{U}(\omega)$  is the free-particle, collisionless propagator and  $L$  is an integral operator, Eq. (22) is an integral equation for  $\hat{U}^\nu(\omega)$ . In our study of problems of this type, we have found that if the functions that appear in the problem are continuous, integrable, and obey some reasonable uniform bounds, then the iteration procedure yields the Neumann series for the associated integral equation and

$$\tilde{U}^\nu(\omega) = \tilde{U}(\omega) + \sum_{n=1}^{\infty} \nu^n [\tilde{U}(\omega)L]^n \tilde{U}(\omega), \quad (27)$$

is the power-series solution for  $\tilde{U}^\nu(\omega)$ , uniformly convergent in the neighborhood of  $\nu=0$ . In this case the iterated  $\epsilon^\nu(\omega)$  is:

$$\epsilon^\nu(\vec{k}, \omega) = 1 - i \frac{\omega_p^2}{k^2} \sum_{n=0}^{\infty} \nu^n T_n(\vec{k}, \omega), \quad (28)$$

$$T_n(\vec{k}, \omega) = \int d^3v \left[ \left( \frac{i}{\omega - \vec{k} \cdot \vec{v}} \right) L \right]^n \left( \frac{i}{\omega - \vec{k} \cdot \vec{v}} \right) \times \vec{k} \cdot \frac{\partial}{\partial \vec{v}} f_0(\vec{v}), \quad (29)$$

where  $f_0$  has been normalized to one and the expression for the free-particle, collisionless propagator has been used:

$$\tilde{U}(\omega) = \int d^3v' \frac{i}{\omega - \vec{k} \cdot \vec{v}'} \delta(\vec{v} - \vec{v}') . \quad (30)$$

For problems of this type, it also follows that Eq. (28) is the power-series solution for  $\epsilon^\nu(\omega)$ , uniformly convergent in the neighborhood of  $\nu=0$ . If Eq. (27) can be summed in closed form and analytically continued for  $\text{Im } \omega < b$ , then the initial-value problem is solved.

## 6. SYMMETRY PROPERTIES

It can be shown that  $\tilde{G}^{\nu}(\vec{k}, \vec{v}, \vec{v}', \omega)^* = \tilde{G}^{\nu}(-\vec{k}, \vec{v}, \vec{v}', -\omega^*)$  with similar formulae for the other  $\vec{k}-\vec{v}-\omega$  quantities discussed in Sections 2 and 3, and that the roots of the collisional dispersion relation,  $\epsilon^{\nu}(\omega)=0$ , obey the relation  $\omega^{\nu}(\vec{k}) = -\omega^{\nu}(-\vec{k})^*$ . The same symmetry properties hold for each of the iterated  $\vec{k}-\vec{v}-\omega$  quantities discussed in Sections 4 and 5 when truncated to the Nth order in  $\nu$ .

## 7. SOME APPLICATIONS

For the free-particle, collisionless propagator, three types of linearized collision operators have been studied: integral, differential, and integral-differential.

As an example of the integral-type collision operator, we studied the simple BGK<sup>2</sup> collision operator, given by

$$L^{\nu} = \nu [-1 + f_0(\vec{v}) \int d^3v'] , \quad (31)$$

where  $f_0$  has been normalized to one. We found that the iterated  $\tilde{U}^{\nu}(\omega)$  and the iterated  $\epsilon^{\nu}(\omega)$  are uniformly convergent power-series solutions in  $\nu$  that agree with expansions of the exact results.

The linear simple Fokker-Planck collision operator is a differential operator and has been studied by Lenard and Bernstein.<sup>3</sup> It is

$$L^{\nu} = \nu \frac{\partial}{\partial \vec{v}} \cdot \left( \vec{v} + v_T^2 \frac{\partial}{\partial \vec{v}} \right) , \quad (32)$$

where  $v_T$  is the electron thermal speed. We found that the iterated  $\tilde{U}^{\nu}(\omega)$  is a divergent, asymptotic expansion as  $\nu \rightarrow 0$  in the sense described above, yet the iterated  $\epsilon^{\nu}(\omega)$  for  $f_0$  Maxwellian is a uniformly convergent power-series solution in  $\nu$  and agrees with the expansion of the closed-form result for  $\epsilon^{\nu}(\omega)$ . For example, the solution to the dispersion relation in the long wavelength limit to order  $\nu$  yields the total damping rate  $\Gamma_k^{\nu}$ , which is the sum of the Landau (collisionless) part  $\gamma_k^{L \rightarrow}$  and the collisional part  $\gamma_k^{\nu \rightarrow}$ , where

$$\gamma_k^{\nu \rightarrow} = -\nu [ 1/2 + 2(k/k_D)^2 + \dots ] . \quad (33)$$

Here  $k_D$  is the Debye wavenumber. Note that these results are obtained without solving a differential equation involving  $L^{\nu}$ .

The linearized Balescu<sup>5</sup> - Lenard<sup>6</sup> collision operator is an integral-differential operator. We have found that the iterated  $\tilde{U}^{\nu}(\omega)$  is a divergent, asymptotic expansion as  $\nu \rightarrow 0$  and we have obtained closed-form results for the iterated  $\epsilon^{\nu}(\omega)$  to the first order in  $\nu$ . Using these results, we have found the solution for the collisional dispersion relation in the long wavelength limit for the linearized Balescu - Lenard collision operator and for  $f_0$  Maxwellian. We obtain the total damping rate  $\Gamma_{\vec{k}}^{\nu}$  which is the sum of the Landau (collisionless) part  $\gamma_{\vec{k}}^L$  and the collisional part  $\gamma_{\vec{k}}^{\nu \rightarrow}$ , where

$$\gamma_{\vec{k}}^{\nu \rightarrow} = -\frac{4}{5} \nu_{ee} \left(\frac{k}{k_D}\right)^2 \left[1 + \frac{69}{14} \left(\frac{k}{k_D}\right)^2 + \dots\right] - \frac{1}{2} \nu_{ei} \left[1 - 2 \left(\frac{k}{k_D}\right)^2 + \dots\right], \quad (34)$$

$$\nu_{ee} = \frac{4}{3} \pi^{1/2} n_0 \frac{q_e^4}{m_e^{1/2} T_e^{3/2}} \ln \Lambda. \quad (35)$$

Here  $\nu_{ei} = 2^{1/2} \nu_{ee}$ ,  $k = |\vec{k}|$ ,  $k_D$  is the Debye wave number,  $n_0$  is the unperturbed density,  $T_e$  is the unperturbed temperature, and  $\ln \Lambda$  is the Coulomb logarithm. Equation (34) is new and was obtained in collaboration with B. Basu. A detailed discussion of its derivation is quite lengthy and will be published elsewhere.

## 8. DISCUSSION

Equations (21) through (23) are the main results of this paper. They give equations for the collisional propagator in terms of the collisionless propagator and the linearized collision operator. An iterative solution of Eq. (22) or (23), given by Eq. (24), which is what we call the expansion method of solution, when substituted into Eqs. (15) through (17) provides a solution of the linearized plasma kinetic equations for a given collision operator. This method of solution as applied to the collisional plasma problem discussed here is new and is particularly useful when a direct solution of the problem cannot be found. For example, the problem of finding the direct solution of the linearized plasma kinetic equations with the Balescu-Lenard collision term and then using it to find the collisional damping of the electrostatic plasma waves appears to be intractable. But by applying the expansion method, we have been able to obtain the collisional damping rate for  $f_0$  Maxwellian to the first order in the plasma parameter, which is given by Eqs. (34) and (35).



## References

1. Landau, L. D. (1946) Zh. Eksp. Teor. Fiz. 16:574 and J. Phys. (USSR) 10:25.
2. Bhatnagar, P. L., Gross, E. P., and Krook, M. (1954) Phys. Rev. 94:511.
3. Lenard, A., and Bernstein, I. B. (1958) Phys. Rev. 112:1456.
4. Dougherty, J. P. (1964) Phys. Fluids 7:1788.
5. Balescu, R. (1960) Phys. Fluids 3:52.
6. Lenard, A. (1960) Ann. Phys. (N. Y.) 3:390.
7. Titchmarsh, E. C. (1948) Introduction to the Theory of Fourier Integrals, Oxford University Press, London, Chapter 1.

END

FILMED

2-30-4

DTIC